

Understanding Hazel

For years, Hazel has faithfully adhered to her prescribed medications, which included digoxin, to help manage her heart failure and atrial fibrillation.

One evening, Hazel started experiencing nausea and mild abdominal pain. Her symptoms worsened over the next two days, and her family grew concerned about her unusual behavior and symptoms. Recognizing the urgency of the situation, they decided to take her to the emergency department.

Hope for Hazel: A DIGIFab Success Story

Presenting Symptoms

- Altered mental status, confusion
- Abdominal pain
- Nausea, vomiting

Current Medications

- Digoxin 0.125 mg QD
- Furosemide 40 mg QD, increased from 20 mg 7 days ago
- Warfarin 2.5 mg QD
- Sacubitril 97 mg/valsartan 103 mg BID

Physical and Neurologic Examinations

- Heart rate 49 bpm
- Other vitals and neurologic examination are normal


Other Examinations

- Laboratory
 - Potassium: 2.1 mEq/L
 - CrCl: 35 mL/min
 - Serum digoxin: 3.0 ng/mL at 6 h post-ingestion
- Electrocardiogram
 - Second-degree AV block

Diagnosis

- Digoxin toxicity

Not an actual patient.



Hazel

Age: 79

Weight: 68 kg

Medical History

- Heart failure with reduced EF
- Atrial fibrillation
- Renal impairment
- Hypertension

See how DIGIFab helped Hazel 

AV, atrioventricular; BID, twice daily; bpm, beats per minute; CrCl, creatinine clearance; EF, ejection fraction; QD, once daily.

INDICATIONS AND USAGE

DIGIFab is indicated for the treatment of patients with life-threatening or potentially life-threatening digoxin toxicity or overdose, including:

DIGIFab[®]
digoxin immune fab (ovine)